

GENERALIZED SUPERSYMMETRIES AND COMPOSITE STRUCTURE IN M -THEORY*

JERZY LUKIERSKI

*Institute for Theoretical Physics, University of Wrocław
pl. M. Borna 9, 50-205 Wrocław, Poland*

Received (received date)

Revised (revised date)

We describe generalized $D = 11$ Poincaré and conformal supersymmetries. The corresponding generalization of twistor and supertwistor framework is outlined with $OSp(1|64)$ superspinors describing BPS preons. The $\frac{k}{32}$ BPS states as composed out of $n = 32 - k$ preons are introduced, and basic ideas concerning BPS preon dynamics is presented. The lecture is based on results obtained by J.A. de Azcarraga, I. Bandos, J.M. Izquierdo and the author¹.

1. Introduction

M -theory has been proposed as a hypothetical quantum theory describing elementary level of matter, which should incorporate and possibly explain various properties of “new string theory” (for review see e.g.²⁻⁴). One of the features of such new theory of fundamental interactions should be the appearance of many extended elementary objects (p -(super)branes, D -(super)branes etc.) related with each other via duality/dimensional reductions net. Such a variety of basic objects in the theory makes sensible a search for some underlying composite structure.

The basic dynamical degrees of freedom in M -theory yet are not known - there were presented only some proposals usually related with $D = 11$ space-time geometry. We postulate that the composite structure of M -theory should be formulated in terms of new degrees of freedom related with new geometry. Because M -theory is supersymmetric, and supersymmetry reveals more elementary nature of spinorial objects, we shall postulate that the basic fundamental geometric structure in M -theory is spinorial.

The only well-known part of the description of M -theory is algebraic. Assuming that M -theory lives in $D = 11$ (this assumption is consistent with description of $D = 11$ SUGRA as the low energy limit of M -theory) we can postulate the following

*Supported by KBN grant 5PO3B05620

basic $D = 11$ M -superalgebra[†]

$$\{Q_r, Q_s\} = Z_{rs}(\Gamma_\mu C)_{rs} P^k + C\Gamma_{[\mu\nu]}C)_{rs} Z^{[\mu\nu]} + (\Gamma_{[\mu_1-\mu_s]}C)_{rs} Z^{[\mu_1\cdots\mu_5]}. \quad (1)$$

where $\mu, \nu = 0, 1, \dots, 10$, $r, s = 1, \dots, 32$. The collection of 528 Abelian generators Z_{rs} ($Z_{rs} = Z_{sr}$) describes the generalized momenta in M -theory. Introducing dual generalized coordinate space

$$X_{rs} = (\Gamma_\mu C)_{rs} X^\mu + (\Gamma_{[\mu\nu]}C)_{rs} X^{[\mu\nu]} + (\Gamma_{[\mu_5-\mu_s]}C)_{rs} X^{[\mu_1\cdots\mu_5]}, \quad (2)$$

we obtain large generalized phase space, with coordinates and positions described by the adjoint representations of $Sp(32)$ algebra.

Let us recall the assumption of Penrose twistor formalism in $D = 4$ ⁸⁻¹¹ that basic spinorial degrees of freedom in twistorial theory of elementary particles are described by N twistors ($i = 1 \dots N$)

$$t_\alpha^{(i)} = \left(\lambda_A^{(i)}, \omega^{(i)\dot{A}} \right), \quad (3)$$

where $\lambda_A^{(i)}, \omega^{(i)\dot{A}}$ ($A = 1, 2$) are the pairs of $D = 4$ Weyl spinors. The following formula for the composite fourmomentum is assumed^{9,11}

$$P_{A\dot{B}} = \sum_{i=1}^N \lambda_A^{(i)} \bar{\lambda}_{\dot{B}}^{(i)}, \quad (4)$$

where $P_{A\dot{B}} = \frac{1}{2} \sigma_{AB}^\mu P_\mu$. We shall propose analogous formula in $D = 11$ for generalized momenta

$$Z_{rs} = \sum_{i=1}^N \lambda_r^{(i)} \lambda_s^{(i)}, \quad (5)$$

where λ_r ($r = 1 \dots 32$) are $D = 11$ real Majorana spinors. In $D = 4$ the twistors (3) are the fundamental representations of the spinorial covering $SU(2, 2)$ of $D = 4$ conformal algebra ($SU(2, 2) = SO(\overline{4, 2})$). In $D = 11$ there exists only minimal conformal spinorial algebra¹²⁻¹⁴ describing the classical real algebra $Sp(64)$, containing $D = 11$ conformal algebra

$$SO(\overline{11, 2}) \subset Sp(64; R). \quad (6)$$

In Sect. 2 we shall consider the generalization of $D = 11$ Poincaré and conformal superalgebras, supersymmetrizing the minimal $D = 11$ conformal spinorial algebra. In Sect. 3 we shall introduce in $D = 11$ the generalization of twistor and super-twistor formalism, with the extensions of Penrose-Ferber relations, which relate $OSp(1|64)$ supertwistor space described by real coordinates ($\xi^2 = 0; R = 1 \dots 65$)

$$T_R = (\lambda_r, \omega^r, \xi), \quad (7)$$

[†]The relation (1) is the standard, minimal M -superalgebra. One can also add arbitrary spin-tensor central charges (see e.g.⁵⁻⁶). The most general case was considered by Sezgin⁷.

with the generalized phase space (X_{rs}, P_{rs}) (see (1-2)). In Sect. 4 we shall describe algebraically $\frac{k}{32}BPS$ states by $n = 32 - k$ superspinors (7) representing $D = 11$ generalized supertwistors. These supertwistorial constituents we shall call BPS preons. It appears that our model geometrically corresponds to new type of Kaluza-Klein theory, with discrete internal extension of space-time coordinates.

2. $D = 11$ Conformal M -(Super)Algebra

Let us observe that the $D = 4$ conformal algebra $(P_\mu, M_{\mu\nu}, D, K_\mu)$ is endowed with the following three - grading structure

$$\begin{array}{ccc} L_1 & L_0 & L_{-1} \\ P_\mu & M_{\mu\nu}, D & K_\mu \end{array} . \quad (8)$$

Grading (8) is determined by the scale dimensions of generators

$$[D, P_\mu] = P_\mu, \quad [D, M_{\mu\nu}] = 0, \quad [D, K_\mu] = -K_\mu \quad (9)$$

and it is easy to see that the conformal algebra (8) has two Poincaré subalgebras: $(P_\mu, M_{\mu\nu})$ and $(K_\mu, M_{\mu\nu})$. For $D = 4$ superconformal algebra $SU(2, 2; 1) = (P_{\mu\nu}, M_{\mu\nu}, D, A, K_\mu; Q_A, \bar{Q}_{\dot{A}}, S_A, \bar{S}_{\dot{A}})$ the three-grading (8) is extended to the following five-grading

$$\begin{array}{ccccc} L_1 & L_{1/2} & L_0 & L_{-1/2} & L_{-1} \\ P_\mu & Q_A, \bar{Q}_{\dot{A}} & M_{\mu\nu}, D, A & S_A, \bar{S}_{\dot{A}} & K_\mu \end{array} , \quad (10)$$

where consistently

$$\begin{aligned} [D, Q_A] &= \frac{1}{2}Q_A, & [D, S_A] &= -\frac{1}{2}S_A, \\ [D, \bar{Q}_{\dot{A}}] &= \frac{1}{2}\bar{Q}_{\dot{A}}, & [D, \bar{S}_{\dot{A}}] &= \frac{1}{2}\bar{S}_{\dot{A}}, \end{aligned} \quad (11)$$

and again $SU(2, 2; 1)$ contains as subalgebras the Poincaré superalgebras $(P_\mu, M_{\mu\nu}, Q_A; \bar{Q}_{\dot{A}})$ and $(K_\mu, M_{\mu\nu}, S_A, \bar{S}_{\dot{A}})$.

The structure of $D = 11$ generalized superconformal algebra, which we call conformal M -superalgebra is quite analogous. The $D = 11$ conformal M -algebra $Sp(64)$ can be in analogy to (8) described by the following three-grading

$$\begin{array}{ccc} L_1 & L_0 & L_{-1} \\ Z_{rs} & R_{rs} & \tilde{Z}_{rs} \end{array} . \quad (12)$$

528 Abelian $GL(32; R)$ 528 Abelian
generators algebra generators

We see that $Sp(64)$ contains two copies of generalized $D = 11$ Poincaré algebras, described by inhomogeneous $Sp(32)$ algebras ($Sp(32; R) \subset GL(32; R)$) with 528 Abelian translation generators.

The superextension of $D = 11$ conformal M -algebra $OSp(1; 64)$ which we call conformal M -superalgebra is described by the following five-grading (see also^{15,16})

$$\begin{array}{ccccc} L_1 & L_{1/2} & L_0 & L_{-1/2} & L_{-1} \\ Z_{rs} & Q_r & R_{rs} & S_r & \tilde{Z}_{rs} \end{array}, \quad (13)$$

where (Q_r, S_r) are the pair of 32-component supercharges, transforming as fundamental representations of $Sp(32)$, with $R_{rs} \subset Sp(32)$ if $R_{rs} = R_{sr}$. The subalgebras spanned by the generators (Q_r, Z_{rs}) and $(S_r, \tilde{Z}_{rs})$ describe two copies of M -superalgebra given by the relations (1).

It should be added that the gradings (12,13) correspond to the grading structure of real Jordanian (super) algebras [17,18].

3. $D = 11$ Supertwistors and Their Relation with Generalized Superspace

Let us recall two basic relations of Penrose twistor theory in $D = 4$ ⁸⁻¹¹

- (i) relation between the generators of Poincaré algebra and twistor components (2)

$$P_{A\dot{B}} = \lambda_A \lambda_{\dot{B}}, \quad (14)$$

$$M_{AB} = \lambda_{(A} \bar{\omega}_{B)}, \quad M_{\dot{A}\dot{B}} = \bar{\lambda}_{(\dot{A}} \omega_{\dot{B})} \quad (15)$$

where $M_{AB} = \frac{1}{2}(\sigma_{\mu\nu})_{AB} M^{\mu\nu}$ and $M_{\dot{A}\dot{B}} = \frac{1}{2}(\tilde{\sigma}_{\mu\nu})_{\dot{A}\dot{B}} M^{\mu\nu\dagger}$. The relations (15) can be extended to all 15 generators of $D = 4$ conformal algebra.

- (ii) Penrose incidence relation between twistor and space-time coordinates

$$\omega^{\dot{A}} = i\lambda_B X^{B\dot{A}} \quad \bar{\omega}^A = -iX^{A\dot{B}} \bar{\lambda}_{\dot{B}} \quad (16)$$

where $X^{B\dot{A}} = (X^{A\dot{B}})^*$ describe four real Minkowski coordinates if the $SU(2, 2)$ twistor norm vanishes

$$(t, t) \equiv i \left(\lambda_A \bar{\omega}^A - \bar{\lambda}_{\dot{A}} \omega^{\dot{A}} \right) = 0. \quad (17)$$

The relations (14-17) can be supersymmetrized. If we introduce the $D = 4$ supertwistor (t_α, η) , which is the fundamental representation of $SU(2, 2; 1)$ with complex Grassmann variable η ($\eta^2 = \bar{\eta}^2 = 0$, $\{\eta, \bar{\eta}\} = 0$), the relations (14-15) has been extended by Ferber¹⁹ to all generators of $D = 4$ superconformal group $SU(2, 2; 1)$.

The Penrose relations (16-17), firstly supersymmetrized in¹⁹ look as follows

$$\omega^{\dot{A}} = i\lambda_B Z^{B\dot{A}} \equiv i\lambda_B \left(X^{B\dot{A}} - i\theta^B \theta^{\dot{A}} \right)$$

[†]We recall that $(\sigma_{\mu\nu})_{AB} = \frac{1}{2i}[(\sigma_\mu)_{A\dot{B}} \tilde{\sigma}_{\nu}^{\dot{B}}{}_{\dot{B}} - (\sigma_\nu)_{A\dot{B}} \tilde{\sigma}_{\mu}^{\dot{B}}{}_{\dot{B}}] = -\frac{i}{2}\epsilon_{\mu\nu\rho\tau}(\sigma^{\rho\tau})_{AB} = [(\tilde{\sigma}_{\mu\nu})_{\dot{B}\dot{A}}]^*$

$$\begin{aligned}\bar{\omega}^A &= i \left(X^{A\dot{B}} + i\theta^A \theta^{\dot{B}} \right) \lambda_B \\ \eta &= \lambda_A \theta^A \quad \bar{\eta} = \bar{\lambda}_{\dot{A}} \bar{\theta}^{\dot{A}}.\end{aligned}\tag{18}$$

For $D = 11$ the generalized twistors and supertwistors are real (see 7) and the real $OSp(1; 64)$ superalgebra ($R, S = 1 \dots 64$)

$$\{Q_R, Q_S\} = R_{RS},\tag{19}$$

can be obtained if we assume that[§]

$$R_{RS} = T_R T_S \quad Q_R = \frac{1}{\sqrt{2}} T_R \xi,\tag{20}$$

where T_R describes $D = 11$ real twistorial quantum phase space ($\eta_{RS} = -\eta_{SR}$ is the $Sp(64)$ antisymmetric metric)

$$[T_R, T_S] = i\eta_{RS},\tag{21}$$

supplemented with trivial one-dimensional Clifford algebra relation $\xi^2 = 1$.

The relations (19) are extended to $D = 11$ as follows:

$$\omega^r = (X^{rs} - i\theta^r \theta^s) \lambda_s \quad \xi = \theta^r \lambda_r.\tag{22}$$

Relations (22) relate the $D = 11$ supertwistor space coordinates (7) with the extended $D = 11$ superspace (X_{rs}, θ_s) , described by 528 bosonic and 32 fermionic coordinates.

4. BPS States in M -Theory and Composites of BPS Preons

The $\frac{k}{32}$ BPS state $|k\rangle$ can be defined as an eigenstate of generalized momenta generators

$$Z_{rs}|k\rangle = z_{rs}|k\rangle,\tag{23}$$

such that $\det z_{rs} = 0$. The number k determines the rank of generalized momenta matrix z_{rs}

$$\frac{k}{32} \text{BPS state: } \{\text{rank } z_{rs} = n = 32 - k; \quad 1 \leq k < 32\}.\tag{24}$$

From (24) follows that the BPS state $|k\rangle$ preserves a fraction $\nu = \frac{k}{32}$ of supersymmetries.

We call BPS preon the hypothetical primary object carrying the following generalized momenta¹

$$Z_{rs} = \lambda_r \lambda_s.\tag{25}$$

[§]By Bott periodicity this realization is related with twistor framework in $D = 3$ (see²⁰), also with real structure. In $D = 5, 6, 7$ one has to use the extension of Penrose framework to quaternionic twistors (see e.g.²¹ for $D = 6$).

The formula (25) corresponds to putting $n = 1$ in the relation (5) and describes the $\frac{31}{32}BPS$ state. More general formula (5) describes the generalized momenta of a system composed out of n BPS preons and it describes (for $1 \leq n \leq 32$) the $\frac{k}{32}BPS$ state (we recall that $k = 32 - n$).

The number $n = 32 - k$ of zero eigenvalues of the matrix z_{rs} determines the number of independent supercharges $Q_r^{(i)}$, annihilating the BPS state $|k\rangle$. These supersymmetries, preserving the BPS state, are called in p -brane theory the κ -transformations. We see that the supersymmetric $D = 11$ single BPS preon dynamics should have 31 κ -symmetries. Recently²² such dynamical superparticle models[¶] with fundamental $OSp(1; 2n)$ superspinor as basic variable has been proposed. It should be recalled here (see e.g.²⁴) that in the standard super p -brane formulations half of the supersymmetries are promoted to κ -transformations, i.e. in $D = 11$ we obtain 16 κ -transformations.

Using the $D = 11$ supertwistor description with the relations (22) and (25) providing a bridge between BPS preons and generalized space-time, we can formulate three different geometric pictures:

- (i) Purely supertwistorial picture, with basic phase space parametrized by BPS preon coordinates $T_R^{(i)}$ (see (7)). The canonical Liouville one-form describing free action is given by the relation

$$\Omega_1 = \sum_{i=1}^n \left(\omega^{(i)r} d\lambda_r^{(i)} + i\xi^{(i)} d\xi^{(i)} \right), \quad (26)$$

which can be supplemented by some algebraic constraints.

- (ii) Mixed geometric picture, with the components $\omega^{(i)}$ expressed by means of the relation (22). One obtains from (26)

$$\Omega_2 = \sum_{i=1}^n \lambda_r^{(i)r} \lambda_s^{(i)r} (dX^{rs} - i\theta^r d\theta^s), \quad (27)$$

- (iii) Generalized space-time picture, with the relation (5) inserted in (27).

$$\Omega_3 = Z_{rs} (dX^{rs} - i\theta^r d\theta^s). \quad (28)$$

The application of these three geometric pictures to the description of $D = 11$ dynamics (for $n > 1$) is under consideration.

5. Final Remarks

We mention here two interesting aspects of the presented approach which deserve further attention;

[¶]For $D = 4, 6$ and 10 see^{22,23}.

(i) geometric confinement of *BPS* preons

Because the space-time coordinates are composed out of preonic degrees of freedom, the $D = 11$ space-time point can be determined only in terms of at least 16 preonic set of spinorial coordinates. This is the $D = 11$ extension of known property of Penrose theory in four dimensions with two twistors needed for the definition of composite Minkowski space-time points.

(ii) internal symmetries

The formula (5) expresses 528 generalized momenta in terms of $32n$ preonic spinorial coordinates $\lambda_r^{(i)}$ ($i = 1, \dots, n$). The internal symmetries can be obtained by interchanging *BPS* preons. For the case $n=16$ corresponding to the choice of $\nu = \frac{1}{2}\text{SUSY}$ one can introduce internal $O(16)$ symmetries, leaving the values of Z_{rs} invariant.

Acknowledgements

The author would like thank prof. Mo-Lin Ge for his warm hospitality at Nankai Symposium in Tianjin.

References

1. I.A. Bandos, J.A. de Azcarraga, J.M. Izquierdo and J. Lukierski *Phys. Rev. Lett.* **86**, 4451 (2001).
2. M.J. Duff, R.R. Khuri and J.X. Lu, *Phys. Rep.* **259**, 213 (1995).
3. P.K. Townsend, *hep-th/9712004*.
4. K.S. Stelle, *hep-th/9803116*.
5. E. Bergshoeff and E. Sezgin, *Phys. Lett.* **B232**, 96 (1989); **B354**, 256 (1995).
6. C. Chrissomalakis, J.A. de Azcarraga, J.M. Izquierdo and J.C. Perez Bueno, *Nucl. Phys.* **B567**, 293 (2000).
7. E. Sezgin, *Phys. Lett.* **B392**, 321 (1997).
8. R. Penrose and M.A.H. Mac Callum, *Phys. Rep.* **6**, 241 (1972).
9. Z. Perjes, *Phys. Rev.* **D11**, 2031 (1975).
10. R. Penrose, *Rep. Math. Phys.* **12**, 65 (1977).
11. L.P. Hugston, "Twistors and Particles", Lecture Notes in Physics, Vol **97** (Springer Verlag, Berlin, 1979).
12. S. Ferrara, *Fortschr. Phys.* **49**, 485 (2001).
13. R. D'Auria, S. Ferrara, M.A. Lledo, V.S. Varadarajan *J. Geom. Phys.* **40**, 101 (2001).
14. R. D'Auria, S. Ferrara, M.A. Lledo, *Lett. Math. Phys.* **57**, 123 (2001).
15. I. Bars, *Phys. Lett.* **B457**, 275 (1999); **B483**, 248 (2000).
16. O. Baerwald and P. West, *Phys. Lett.* **B476**, 157 (2000).
17. M. Cederwall, *Phys. Lett.* **B115**, 389 (1988).
18. M. Gunaydin, *Mod. Phys. Lett.* **A15**, 1407 (1993).
19. A. Ferber, *Nucl. Phys.* **B132**, 55 (1978).
20. W. Heidenreich and J. Lukierski, *Mod. Phys. Lett.* **A5**, 439 (1990).
21. J. Lukierski and A. Nowicki, *Mod. Phys. Lett.* **A6**, 189 (1991).
22. I. Bandos, J. Lukierski and D. Sorokin, *Phys. Rev.* **D61**, 045002 (2000).
23. I. Bandos, J. Lukierski, Ch. Preitschopf and D. Sorokin, *Phys. Rev.* **D61**, 065009 (2000).
24. D. Sorokin, *Phys.Rep.* **329**, 1 (2000).